

# FujiMite® XLO on HOPS

Control two-spotted spider mite with  
FujiMite XLO miticide/insecticide.

**New  
Low Odor  
Formulation**

**FujiMite® XLO miticide/insecticide provides fast knockdown and lasting control of two-spotted spider mites in hops.**

## Key Benefits

- Active on all developmental stages of mites – larvae, nymphs, and adults
- Stops mite feeding within hours of application
- Provides lasting control by inhibiting molting on immatures
- Easy on most beneficial insects
- Excellent fit for resistance management
- Apply mid-to-late season for excellent mite knockdown and lasting control

## Recommended Use Rates

- 3 pints per acre
- Spray concentrations above 100 ppm are recommended with full coverage high volume spray equipment.

## Use Information

- Apply FujiMite XLO before mite populations exceed 5 mites per leaf.
- FujiMite XLO is a contact miticide. Proper application coverage is important for maximizing control of mites.
- Leaf yellowing may occur if FujiMite XLO is combined with spray oil in excess of 1% of the spray volume.
- Reentry interval – 12 hours
- Preharvest interval – 15 days
- Group 21A insecticide

Aaron Gale  
**Technical Sales Representative**  
509-679-9756



### Mite Fact:

Individual females can lay up to 20 eggs per day. Development from egg to adult takes 7 to 10 days. Normally, about nine generations of mites develop on hops during the summer months.

**FujiMite® XLO**  
MITICIDE/INSECTICIDE

**NICHINO**  
AMERICA