



For Fast Knockdown of Mites with Added Benefit of Leafhopper Control

GRAPE

FujiMite® XLO miticide/insecticide provides control on a wide range of mites in table, raisin, and wine grape varieties. FujiMite XLO will also control leafhoppers and provide contact control of mealybugs which may be present at the time of application.

Key Benefits

- Active on all motile stages of mites - larvae, nymphs, and adults and contact activity on existing eggs
- Fast knockdown of mites
- Long-lasting control by inhibiting molting of immatures
- Unique mode-of-action stops mite feeding damage immediately and oviposition within hours after application
- Provides control of leafhoppers and mealybugs; suppresses powdery mildew

Key Mites Controlled


Pacific spider mite, Willamette spider mite, Two-spotted spider mite

Use Recommendations


- Rates
 - Willamette spider mite: 1.5 - 2.0 pts/A
 - All other mites and mealybugs: 2.0 pts/A
 - Leafhoppers: 1.0 - 2.0 pts/A
- Apply by ground using a minimum of 50 gallons of water per acre.
- Application speed for optimal coverage is LESS than 2 mph.

Use Information

- FujiMite XLO is a contact miticide. Proper application coverage is critical for maximizing mite control.
- Reentry interval – 12 hours
- Preharvest interval – 14 days
- Group 21A Insecticide




FujiMite XLO Efficacy on Pacific Spider Mite on Grape in Kerman, CA 2011



Average pretreatment counts were 327 mites per leaf
 Spray volume: 100 GPA
 NIS 0.25% v/v

FujiMite XLO Efficacy on Pacific Spider Mite on Grape in Kerman, CA 2013



Average pretreatment counts were 535 mites per leaf
 Spray volume: 100 GPA
 NIS 0.25% v/v

