

Convoy® fungicide protects rice from sheath blight caused by strobilurin resistant and susceptible *Rhizoctonia* strains. Flutolanil, the active ingredient in Convoy, is a SDHI fungicide that provides both curative and preventative control of sheath blight.

### Key Benefits

- Provides effective control of sheath blight in areas where strobilurin resistant *Rhizoctonia* strains are present.
- Non-strobilurin mode of action – to control or prevent development of resistant sheath blight, FRAC Group 7
- Dependable control of sheath blight through preventative and curative action
- Systemic activity, providing protection throughout the entire sheath
- Flexible application program – may be used in a single or two-spray program
- Excellent yield and quality results
- 0 day plantback interval for cotton and soybeans; 30 day plantback interval for wheat

### Use Recommendations


- Single Application Program
  - Rate: 32 fl oz/acre
  - Apply early-boot to boot
- Two Application Program
  - Rate: 16 fl oz/acre
  - Apply early boot and late boot

### Use Information

- Convoy is only active on sheath blight in rice. For control of additional diseases, another fungicide should be tank mixed with Convoy.
- A minimum of 5 gallons of water must be used when applying Convoy by air.



**Strobilurin Resistant Sheath Blight Severity, LA\***



Applications made at boot


\*Trial conducted in a commercial rice field with a history of strobilurin resistant *Rhizoctonia solani*

Dr. Don Groth, LSU AgCenter, Mowata, LA, 2014



# For Control of Sheath Blight

## Strobilurin Resistant Sheath Blight Incidence, LA\*




Applications made at boot

\*Trial conducted in a commercial rice field with a history of strobilurin resistant *Rhizoctonia solani*

Dr. Grady Coburn, Pest Management Enterprises, LLC, Mowata, LA, 2014

## Susceptible Sheath Blight Severity, TX\*




Applications made at boot

\*strobilurin susceptible strain of *Rhizoctonia solani*

Dr. Young-Ki Jo, Texas AgriLife Extension Service, Beaumont, TX, 2014

## Susceptible Sheath Blight Yield, TX\*



Applications made at boot

\*strobilurin susceptible strain of *Rhizoctonia solani*

Dr. Young-Ki Jo, Texas AgriLife Extension Service, Beaumont, TX, 2014

