

**Low Odor
Formulation**

STRAWBERRY

For Control of Damaging Mites

FujiMite® XLO miticide/insecticide provides excellent control on a wide range of mites in strawberries. The active ingredient, *fenpyroximate*, stops mite feeding immediately and oviposition within hours of application.

Key Benefits

- Controls all motile stages of mites – larvae, nymphs, adults
- Fast knockdown of mites and rapid cessation of feeding
- Lasting control by inhibiting molting of immatures
- Provides benefit of whitefly suppression
- Classified by the EPA as reduced risk
- Excellent fit for resistance management – IRAC Group 21A

Key Mites Controlled

Two-spotted spider mite	Pacific spider mite	Strawberry spider mite	Carmine spider mite	Cyclamen mite
✓	✓	✓	✓	✓

Recommended Use Rates

- 2 pints/acre
- Minimum spray volume – 25 gallons of water per acre

Use Information


- FujiMite XLO is a contact miticide. Proper application coverage is important for maximizing the control of mites.
- Reentry interval - 12 hrs
- Preharvest interval – 1 day
- Predatory Mite Release – If predatory mites are to be released after a FujiMite XLO application, please wait approximately two weeks.



Two-spotted Spider Mite



Carmine Spider Mite



Cyclamen Mite

